

# Endothelial Integrity, Soluble Adhesion Molecules and Platelet Markers in Type 1 Diabetes Mellitus

A.D. Blann\*, G.Y.H. Lip

Haemostasis, Thrombosis and Vascular Biology Unit, University  
Department of Medicine, City Hospital, Dudley Road, Birmingham, UK

Recent developments in cell biology have identified new areas of direct relevance to the pathogenesis of Type 1 (insulin-dependent) diabetes mellitus and its complications. Endothelial damage is well recognized in diabetes—endothelial cell markers von Willebrand factor, soluble E-selectin, and soluble thrombomodulin are providing further evidence of the relationship between activation and damage to the vasculature and clinical disease in this condition. Cell surface bound adhesion molecules may also have a role in the development of atherosclerosis in patients with diabetes but the importance of the soluble forms of these molecules, such as intercellular adhesion molecule-1, is unclear. Evidence of platelet dysfunction has long been acknowledged in diabetes and new data are discussed. It is likely that a greater appreciation of the intimate interactions between endothelial integrity, adhesion molecules and platelets in Type 1 diabetes mellitus will provide a greater understanding of the risk of cardiovascular disease and stroke in patients with this disorder. © 1998 John Wiley & Sons, Ltd.

*Diabet. Med.* 15: 634–642 (1998)

**KEY WORDS** Type 1 diabetes mellitus; endothelial cells; platelets; Soluble adhesion molecules

Received 11 November 1997; revised 7 January 1998; accepted 28 February 1998

## Introduction

Subjects with Type 1 (insulin-dependent) diabetes mellitus (DM) are at increased risk of premature mortality and morbidity, mostly through the development of atherothrombotic vascular diseases such as ischaemic heart disease, peripheral arterial occlusive disease and stroke, and other related clinical complications such as retinopathy, neuropathy and nephropathy. The precise pathogenic mechanisms for these changes are unclear but may involve alterations in several physiological systems, including those of haemostasis, thrombosis, and vascular biology and especially in the correct functioning of the endothelial cell and the platelet.<sup>1–5</sup> Many of these changes are also important in, and relevant to, the pathogenesis of Type 2 (non-insulin-dependent) diabetes mellitus and its complications, but as this has been reviewed in a recent issue of *Diabetic Medicine*,<sup>6</sup> and elsewhere (e.g. Schneider and Sobel<sup>7</sup>), this review concentrates upon Type 1 DM. Although information derived from animal work has also provided useful insights, we primarily consider data collected in man.

## Methodology

The ovid medicine database (1990–1997) was searched, using the key words: insulin-dependent diabetes mellitus, endothelial cells, platelets, soluble adhesion molecules. A personal assessment was made of the relevance of each study found to the topic of this review.

## The Endothelium

The importance of the endothelium in cardiovascular pathophysiology is well established, and is discussed in several excellent reviews.<sup>6,8–10</sup> The functioning of this organ may be assessed invasively (e.g. by the perfusion of organs with pharmacologically active agents such as acetylcholine), non-invasively (e.g. by using the hyperaemic (flow mediated dilatation) response) and with the use of plasma markers. Vascular biologists have at their disposal a limited number of these specific plasma markers which reflect endothelial cell physiology and which therefore may be used as tools to dissect the disease process.<sup>9,11</sup>

### von Willebrand factor

More is known of the physiological (e.g. in exercise) and pathological (e.g. in hypertension) relationships of this molecule than any other marker of endothelial cell damage or dysfunction. Boneu and colleagues were

\*Correspondence to: Dr A.D. Blann, Haemostasis, Thrombosis and Vascular Biology Unit, University Department of Medicine, City Hospital, Dudley Road, Birmingham B18 7QH, UK. E-mail: blanna@novell5.bham.ac.uk

among the first to note raised levels in diabetes mellitus although it is unclear whether or not their subjects had Type 1 or Type 2 disease.<sup>12</sup> With even higher levels among patients whose disease was complicated by atherosclerosis, they predicted that raised von Willebrand factor was the result of endothelial damage, a notion now widely accepted.

Tissue culture studies indicate that high glucose levels can cause release of von Willebrand factor from its intracellular storage granule, the Weibel-Palade body, within endothelial cells.<sup>13</sup> Greaves and colleagues showed that resolution of diabetic ketoacidosis may be accompanied by a significant reduction in raised levels of von Willebrand factor and suggested that this may be a mechanism for endothelial damage (Figure 1).<sup>14</sup> Leurs *et al.*<sup>15</sup> found raised von Willebrand factor in Type 1 DM patients, which failed to show a clear relationship with HbA<sub>1c</sub>. Many groups have shown that increased levels of this marker have a positive relationship with both retinopathy and nephropathy in Type 1 DM, and correlations with urinary albumin excretion are commonly described.<sup>16–21</sup> However, Myrup *et al.* were unable to confirm this relationship, instead finding low HDL-cholesterol<sup>22</sup> and raised sialic acid<sup>23</sup> to have stronger associations with microalbuminuria. Knobl *et al.*<sup>24</sup> found a correlation between von Willebrand factor and albuminuria in Type 1, but not Type 2, DM patients. However, raised von Willebrand factor is not restricted to patients with retinopathy or nephropathy: Plater *et al.*<sup>25</sup> showed that raised levels are associated with deteriorating nerve function, independent of glycaemic control.

The association between von Willebrand factor, endothelial cell injury and Type 1 DM may be extended to include a (possibly causative) relationship with increased

systolic blood pressure and oxidant injury, but a relationship with insulin levels is unproven.<sup>26–28</sup> Notably, Stehouwer *et al.*<sup>29</sup> reported that an increase in von Willebrand factor preceded the occurrence of microalbuminuria (but not retinopathy) by approximately 3 years, providing evidence of its value as a potentially useful predictive clinical tool, a finding later confirmed elsewhere.<sup>23</sup> However, Deckert and colleagues showed increased urinary albumin excretion, itself a marker of silent myocardial ischaemia in Type 1 DM,<sup>30,31</sup> to be a better predictor of disease progression than increased levels of von Willebrand factor.<sup>32</sup> However, in this case it cannot be said that the raised von Willebrand factor reflects damage only to the endothelium of the kidney: it is likely to represent a global score of vascular function and is generally unlikely to be a marker of damage to any one specific vascular bed. Hence both raised von Willebrand factor and microalbuminuria would appear to be the result of the same (possibly diabetic) disease process. A similar argument can be applied to the relationship between retinopathy and increased von Willebrand factor: one presumes that the latter is marking endothelial dysfunction, and one aspect of this is damage to the retinal vessels. Increased von Willebrand factor may therefore be the common laboratory measurement of the various mechanisms (e.g. cigarette smoking, free radicals, oxidised lipids) which may bring about endothelial cell injury in a variety of diseases. One consequence of any of these mechanisms may be microalbuminuria, another may be hypertension. It is also possible that high levels of von Willebrand factor take an active part in the disease process by promoting thrombus formation, and it is notable that high levels of this factor predict a poor outcome (e.g. myocardial




Figure 1. Possible mechanisms to explain the relationship between diabetes and atherosclerosis. Text in parentheses refers to potential participation (or not) in the progression of disease

infarction, stroke, death) in cardiovascular disease.<sup>33,34</sup> We would presume, therefore that the predictive power of this molecule for cardiovascular disease progression is equal in both Type 1 and Type 2 DM, but this needs confirmation in long-term outcome studies.

Patients with microalbuminuria often also have significant elevations of plasma fibrinogen, triglycerides, and LDL-cholesterol, and lower levels of HDL-cholesterol, when compared to controls.<sup>35</sup> Therefore both endothelial and lipid abnormalities add significantly to the complex risk factor profile for cardiovascular disease in Type 1 DM patients.

## Soluble Thrombomodulin

Thrombomodulin (CD141) is a proteoglycan component of the endothelial cell membrane with anticoagulant properties. It binds thrombin and converts it into an enzyme that activates the protein C pathway. Consequently, the appearance of a fractionated, soluble form in the plasma implies (by its presumed loss from the endothelial cell surface) not only vascular injury but also a change in haemostasis towards coagulopathy.<sup>36</sup>

Two groups have been unable to find raised levels of this molecule in diabetic subjects with normal albumin excretion rates, but high levels have been found in patients with microalbuminuria and in the presence of clinical nephropathy.<sup>37,38</sup> Gruden *et al.*<sup>39</sup> have also reported higher soluble thrombomodulin in 12 Type 1 DM patients with microalbuminuria when compared to 12 patients with normoalbuminuria. They also found a correlation between soluble thrombomodulin and diastolic blood pressure, a finding similar to that of Gabat *et al.*, who also found that soluble thrombomodulin was more prominent in a mixed group (i.e. Types 1 and 2 DM) of hypertensive patients compared to normotensive patients.<sup>40</sup> Leurs *et al.* were initially unable to find raised soluble thrombomodulin, or a clear relationship with HbA<sub>1c</sub> in 25 Type 1 DM patients free of microvascular complications such as retinopathy compared to 21 healthy controls,<sup>15</sup> but later<sup>41</sup> correlated soluble thrombomodulin with microalbuminuria in a study of 46 patients and 18 healthy controls. Seigneur *et al.*<sup>42</sup> reported the highest levels in diabetic patients with proliferative retinopathy, although it is unclear whether or not their patients had Type 1 or Type 2 DM.

Consequently, there is still considerable scope for research into soluble thrombomodulin in diabetes, especially in outcome and intervention studies. It will be particularly interesting to discover if soluble thrombomodulin levels differ significantly between Types 1 and 2 disease, and if it has more to offer than von Willebrand factor in predicting those patients at increased risk of additional disease. In this respect, for example, McLaren *et al.* reported a preliminary study of raised soluble thrombomodulin, but normal von Willebrand factor, in young (median age 15 years) Type 1 diabetic subjects, suggesting the former is a more sensitive

marker of endothelial dysfunction.<sup>43</sup> Hence soluble thrombomodulin represents another opportunity to study vascular function, but more extensive clinical data are required, especially from large studies. There are currently very few follow-up data for soluble thrombomodulin in diabetes, which is eagerly sought as high levels also imply endothelial damage and a procoagulant profile.

## Soluble E-selectin

Increased levels of E-selectin (CD62E) appear at the surface of the immunologically activated endothelium, and increased soluble levels also appear in tissue culture supernatants when endothelial cells are co-cultured with cytokines and in the plasma of subjects with certain diseases such as septic shock.<sup>44,45</sup> However, as for soluble thrombomodulin, the definitive clinical experiments that clearly show, in large numbers, a relationship between soluble E-selectin and either Type 1 or Type 2 DM are still awaited. Gearing and colleagues were among the first to find increased levels of soluble E-selectin in (unspecified) diabetes,<sup>45</sup> and several groups have extended this work by showing raised levels in both Types 1 and 2 DM<sup>46,47</sup> although not all workers have agreed with this observation.<sup>48</sup> The latter group was also unable to find increased levels among patients with microangiopathy.

We have been unable to find increased soluble E-selectin among patients with non-diabetic atherosclerosis.<sup>49</sup> It is therefore notable that Kirk and colleagues have reported raised levels in diabetic relative to non-diabetic atherosclerosis and to healthy controls.<sup>50</sup> This suggests that raised soluble E-selectin may have a specific relationship to diabetic vascular biology and not necessarily to atherosclerosis *per se*. Preliminary data from the same group indicates that dietary supplementation with essential fatty acids is effective in reducing increased levels of soluble E-selectin in young people with Type 1 DM.<sup>51</sup> This suggests that it may be possible to restore vascular function with diet manipulation, and so represents a new initiative.

## Endothelin

Endothelin is a potent vasoconstrictor derived from the endothelium that acts directly on smooth muscle cells. Its presence in the plasma is presumed to be a spillover from this activity at the interface between the two cell types. Lehman *et al.*<sup>52</sup> hypothesized that plasma endothelin may be an additional marker for arterial vascular disease and speculated that it might be a contributor to atherogenesis or may merely be released from damaged endothelial cells. Takahashi and colleagues were among the first to describe increased levels of plasma endothelin-1 among patients with both Type 1 DM and Type 2 DM.<sup>53</sup> Collier *et al.* have subsequently shown these increased plasma levels are related directly to albuminuria in Type 1 DM<sup>54</sup> while Kawamura *et al.*

have described raised plasma levels in Type 2 DM patients with retinopathy.<sup>55</sup> Curiously, however, levels seem to be *lower* in children and adolescents with Type 1 DM than in their healthy siblings.<sup>56</sup> Smulders *et al.* also found lower plasma levels of endothelin in 10 Type 1 DM adults compared to 10 healthy controls.<sup>57</sup>

Properzi *et al.*<sup>58</sup> examined endothelin in the endothelium of skin biopsies, using immunohistology and electronmicroscopy. They found levels of tissue endothelin-1 to be higher (present in 82 % of cells in the biopsies) in patients with Type 1 DM of less than 10 years duration, than in healthy controls, where only 50 % of cells were stained. However, only 17 % of cells from patients with long standing diabetes were reactive, and the percentage of biopsies showing positive endothelin-1 staining was lower in patients with retinopathy than in patients without retinopathy. No differences in staining for von Willebrand factor were observed. It has been suggested that measurement of plasma endothelin may provide some useful information, especially as a vascular marker,<sup>59,60</sup> and a thorough comparison with other endothelial markers such as von Willebrand factor would be valuable. However, Gruden *et al.*<sup>61</sup> have been unable to find raised plasma endothelin in 13 microalbuminuria patients compared to 13 normoalbuminuric patients.

Therefore at present there seems to be little consensus regarding a role for plasma endothelin in Type 1 DM. The likelihood of false positives and false negatives in the small studies described can only be addressed by studies in large (>100) numbers of patients with age- and sex-matched controls. The relationship between plasma endothelin and insulin also needs to be fully explored.

### Other Endothelial Products

Vascular endothelial cells synthesize an inhibitor of tissue factor (a co-factor for coagulation factor VIIa) called tissue factor pathway inhibitor, or TFPI, and may be its principal source.<sup>62–64</sup> It is unclear whether or not raised levels of this anticoagulant reflect endothelial cell damage, but increased TFPI (defined by a chromogenic, not a clotting, assay) is found in Type 1 DM, correlating with glycated haemoglobin but not with increased von Willebrand factor.<sup>15</sup> A subsequent report from the same group<sup>41</sup> showed higher levels in patients with microalbuminuria compared to patients with retinopathy or patients without microangiopathic complications. Yokoyama *et al.*<sup>65</sup> found that plasma TFPI correlated with urinary albumin excretion (which may be considered a surrogate marker of vascular damage) in 59 patients with Type 1 DM.

Changes in levels of other molecules likely to be important in thrombosis and haemostasis have also been reported. Leurs *et al.*<sup>15</sup> found low levels of tissue plasminogen activator (tPA) in diabetes, but no change in levels of its inhibitor (i.e. plasminogen activator

inhibitor (PAI)). Skrha *et al.*<sup>66</sup> found raised tPA regardless of degree of retinopathy, and Gruden *et al.*<sup>61</sup> found tPA to be uninfluenced by albumin excretion rate, although this group found PAI to be higher in subjects with microalbuminuria. However, the precise value and status of tPA and PAI as specific endothelial cell markers are unclear. For example, they can be measured by immunological or functional methods, and may form a (possibly inactive) complex *in vivo* and/or *in vitro*. Furthermore, some may derive from non-endothelial sources such as platelets.<sup>67–70</sup> An example of the difficulty in interpretation of levels of tPA is the paradox that smoking, widely believed to be pathogenic towards the endothelium, the platelet, and to increase soluble coagulation factors, increases the activity of plasma tPA,<sup>71</sup> which should promote clot dissolution, and so reduce the risk of thrombosis.

### Soluble Adhesion Molecules

In addition to endothelial cell soluble E-selectin already described, several other soluble adhesion molecules have been studied in Type 1 DM: these are intercellular adhesion molecule-1 (ICAM-1, CD106), vascular cell adhesion molecule-1 (VCAM-1, CD54), soluble L-selectin (CD62L), and soluble P-selectin (CD62P). The first two are both members of the immunoglobulin gene superfamily, the last two will be discussed in the section on platelets which follows. Raised soluble L-selectin has been described in Type 1 DM,<sup>72</sup> but there is little other information about this molecule in diabetes which can, in other circumstances, reach levels high enough to inhibit cell to cell adhesion.<sup>75</sup> Least is known about the most recently characterized, platelet–endothelial cell adhesion molecule-1 (PECAM-1, CD31).<sup>73</sup> Levels do not seem to be altered in atherosclerosis<sup>74</sup> and a preliminary report found levels in diabetes to be no different from those in healthy controls.<sup>76</sup> For more comprehensive recent reviews on the cell biology of adhesion molecules (such as their involvement in cell rolling and adhesion) see Carter and Grant,<sup>6</sup> Jang *et al.*,<sup>77</sup> Kansas,<sup>78</sup> and Frenette and Wagner.<sup>79</sup>

#### Soluble ICAM-1

This adhesion molecule is a transmembrane glycoprotein that binds to members of the beta 2 integrin family; leucocyte function associated antigen (CD11a/CD18) and Mac-1 (CD11b/CD18), and to the sialoprotein CD43.<sup>80</sup> Gearing *et al.* were among the first to describe raised plasma levels in diabetes,<sup>45</sup> and Lampeter *et al.* reported increased levels in 4 out of 14 patients with recent onset Type 1 DM and in first degree relatives with a genetic predisposition to the disease.<sup>72</sup> They postulated from their subsequent *in vitro* data that soluble ICAM-1 may downregulate inflammatory responses, providing a novel opportunity to intervene in the disease process.<sup>81</sup> Interestingly, therapy with monoclonal anti-



bodies to ICAM-1 prevents graft rejection and Type 1 DM recurrence in a rat model.<sup>82</sup> Other animal work has led Yagi *et al.* to suggest that ICAM-1 plays an important role in the pathogenesis of Type 1 DM via involvement in the extravasation of lymphocytes from the plasma into the inflamed pancreas.<sup>83</sup> Whether or not soluble ICAM-1 modulates this process in humans remains to be proven.

However, an alternative view is suggested by the data of Accardo-Palumbo *et al.*,<sup>84</sup> who found higher soluble ICAM-1 in patients who were positive for antibodies to myeloperoxidase. They postulated that damage to the endothelium (as suggested by raised soluble ICAM-1) is due to chronic neutrophil activation by antibodies to myeloperoxidase. Both these avenues are worthy of additional study, especially as Fasching *et al.*<sup>48</sup> have correlated raised levels of soluble ICAM-1 with the presence of microangiopathy, although Cominacini *et al.* have not been able to find raised levels of ICAM-1 in Type 1 DM.<sup>47</sup>

Esser and colleagues have taken a novel approach. They reported ICAM-1 to be raised by 13 % in the plasma of Type 1 DM patients with proliferative diabetic retinopathy (PDR), but by only 2 % in the plasma of Type 2 DM patients with PDR.<sup>85</sup> However, they also measured levels in vitreous humour from the patients, finding levels to be about 1 % of those in healthy control plasma. In the Type 1 DM patients with PRD, levels were 5.8 % of those in the plasma, while in Type 2 disease, levels were 7.6 % of paired plasma levels. The highest levels relative to plasma were in idiopathic (8.9 %) and in traumatic PDR (10.4 %). They conclude that local ICAM-1 production, possibly from macrophages, may be of importance in the early phase of retinal disease by enhancing immune activation and inflammation.

### Soluble VCAM-1

This molecule, also a member of the immunoglobulin gene superfamily, is found on endothelial cells and dendritic cells, and weakly raised levels in diabetes have been reported.<sup>45</sup> Fasching *et al.* also described higher levels in 37 patients with Type 1 DM but no microangiopathy than in 70 healthy controls. However, levels were even higher among 33 patients who had microangiopathy compared to the Type 1 patients who were free of microangiopathy.<sup>48</sup> However, as for soluble ICAM-1, Cominacini *et al.*<sup>47</sup> did not find levels of soluble VCAM-1 to be higher in 18 patients with Type 1 DM compared to 20 healthy controls. These discrepancies clearly need to be resolved in larger studies, as they may have implications for vascular biology and the possible use of antibody therapy directed towards the cell surface from of the adhesion molecule. For example, administration of monoclonal antibodies to VCAM-1 inhibited the development of insulinitis and diabetes in a mouse model of Type 1 DM.<sup>86</sup>

A recent exciting report has been the preliminary data

from Abeygunaratne *et al.*<sup>87</sup> They showed that increased soluble ICAM-1, but not soluble VCAM-1 or soluble E-selectin, predicted the development of vascular complications in 11 of 36 people (17 with Type 1 DM, 19 with Type 2 DM) followed up for 5 years. The full report of this study will provide evidence of an additional tool to discover those subjects at risk of additional disease.

### Platelet Markers

Platelets have long been implicated in the pathogenesis of the vascular disease that accompanies Type 1 DM,<sup>88</sup> reviewed in Colwell *et al.*<sup>4,89</sup> The alpha granule component, beta thromboglobulin (Figure 1), has been widely used to look for inappropriate platelet activity in Type 1 DM and its complications.<sup>90–95</sup> Patrick *et al.*<sup>90</sup> concluded that urinary beta thromboglobulin is not sensitive enough to be useful in the detection of early renal disease, despite raised levels being found in diabetes. Rak *et al.*<sup>93</sup> found that plasma levels closely correlated with the progression to angiopathic disease, although most of their cohort had Type 2 DM and Bayraktar *et al.*<sup>94</sup> were unable to find raised levels of beta thromboglobulin in 30 patients with Type 1 DM compared to 15 controls. More recently Huszka and colleagues<sup>95</sup> reported levels of  $169.1 \pm 44.6 \text{ ng ml}^{-1}$  (mean  $\pm$  SD) in 15 patients with Type 1 DM,  $65.2 \pm 18.4 \text{ ng ml}^{-1}$  in 35 patients with Type 2 DM, compared to their non-diabetic reference range of 15–35  $\text{ng ml}^{-1}$ . The differences in these values would appear to be significant, but no statistical analyses were presented. There is a further report that subjects with Type 1 DM have raised beta thromboglobulin<sup>6</sup> and the longitudinal study has shown the marker to be a significant predictor of disease progression.<sup>96</sup> Despite this interest, it is by no means clear that uncomplicated Type 1 DM is associated with platelet activation.

Soluble P-selectin (CD62P) has recently been proposed as a new marker of platelet activation.<sup>97,98</sup> Being shed from the platelet membrane, this adhesion molecule provides an additional perspective to the matrix alpha granule component beta thromboglobulin. Raised levels of both markers are found in non-diabetic atherosclerosis<sup>99</sup> and increased soluble P-selectin has been reported in Type 1 DM.<sup>100</sup> However, it is not yet known whether there are differences according to retinopathy or nephropathy and a good comparison with beta thromboglobulin would provide much needed data. More recently, Rauch *et al.* have used membrane bound P-selectin to provide preliminary data indicating increased numbers of circulating activated platelet in Type 1 DM patients relative to healthy controls.<sup>101</sup> They interpreted lower levels of such activated cells in patients with microangiopathy as evidence of increased consumption. Conversely, Tschoepe *et al.* found higher numbers of platelets bearing P-selectin in prediabetic subjects, concluding that these subjects have platelet activation which may represent an antecedent, potentially pathogenic feature of Type 1 DM.<sup>102</sup>

## Conclusions and Speculations

Physiological techniques for assessing endothelial function in diabetes (such as invasive, pharmacological methods, and the non-invasive flow mediated dilatation studies,<sup>103</sup> are impractical for large epidemiological studies of the aetiopathogenesis of the complications of Type 1 DM. Several plasma molecules, markers of endothelial (dys)function, are available which may be useful in extending our knowledge of the pathophysiology of diabetes. For some, interesting preliminary data have been published but they are often derived from small numbers of subjects and/or from studies of cross-sectional design. In others the lack of a definition of the precise type of diabetes of the patients has lead to difficulties in interpretation. A further difficulty with the interpretation of increased levels of many of these markers (such as soluble ICAM-1) is that they may simply relate to the disturbances in autoimmunity and not necessarily to the insulin/glycosylation aspects of the disease.

Adequate endothelial cell functioning is as crucial in Type 1 DM as it is in other conditions.<sup>9,104</sup> In this respect, there is little doubt that von Willebrand factor has most to offer as a marker of this process. As for platelets, although both beta thromboglobulin and soluble P-selectin are increased, so only the former has been described as a significant predictor of disease progression.<sup>96</sup> Although unlikely to be of value as diagnostic aids, students of the endothelium and platelet will be interested to learn if any of the markers behave differently in the two types of diabetes as this may provide clues to the interaction between the disease processes and the biology of the cell concerned.

Whether or not soluble adhesion molecules are simple markers of various physiological and pathological processes remains to be proven. They may perhaps help provide clinical guidance as to those patients at risk of the development of cardiovascular complications, and may prove to have relevance to cell/cell adhesion *in vivo*. It seems likely that a large fraction of the levels of certain plasma markers, such as tPA and soluble ICAM-1, arises from the endothelium. However, this may be of little interest to the clinician unless they can provide useful information about current disease activity and/or the risk of developing complications. It is possible that the increasing interest in the platelet and endothelium as likely participants in pathogenesis<sup>3,4,105–107</sup> may translate into the desire to direct treatment in this direction. Anti-platelet (e.g. aspirin) and anticoagulant (e.g. Warfarin) drugs for the former are established, but for the endothelium, only antihypertensives (such as ACE inhibitors) may directly act on the endothelium, and as they are common in clinical practice they may represent a possible growth area for those interested in retaining and improving vascular integrity.

## Acknowledgements

We thank S. Jones and C. Stehouwer for constructive critique of the manuscript.

## References

- Steiner G. Diabetes and atherosclerosis. *Diabetes* 1981; **30** (suppl 2): 1–7.
- Morrish NJ, Stevens LK, Head J, Jarrett RJ, Keen H. A prospective study of mortality among middle aged diabetic patients (the London cohort of the WHO multinational study of vascular disease in diabetics). I: Causes and death rates. *Diabetologia* 1990; **33**: 538–541.
- Ross R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature* 1993; **362**: 801–809.
- Colwell JA, Winocour PD, Halushka PV. Do platelets have anything to do with diabetic microvascular diseases? *Diabetes* 1983; **32** (suppl 2): 14–19.
- Vukovich TC, Scherthaner G, Knobi PN, Hay U. The effect of near normoglycaemic control on plasma factor VIII/von Willebrand factor and fibrin degradation products in insulin-dependent diabetic patients. *J Clin Endocrinol Metabol* 1989; **69**: 84–89.
- Carter AM, Grant P. Vascular homeostasis, adhesion molecules and macrovascular disease in non-insulin dependent diabetes mellitus. *Diabetic Med* 1997; **14**: 423–432.
- Schneider DJ, Sobel BE. Determinants of coronary vascular disease in patients with type II diabetes mellitus and their therapeutic complications. *Clin Cardiol* 1997; **20**: 433–440.
- Davies MG, Hagen P-O. The vascular endothelium. A new horizon. *Annals Surg* 1993; **218**: 593–609.
- Poston L, Taylor PD. Endothelium-mediated vascular function in insulin dependent diabetes mellitus. *Clin Sci* 1995; **88**: 245–255.
- Luscher TF. The endothelium and cardiovascular disease—a complex relation. *N Engl J Med* 1994; **330**: 1081–1083.
- Pearson JD. Markers of endothelial perturbation and damage. *Br J Rheumatol* 1993; **32**: 651–652.
- Boneu B, Abbai M, Plante J, Bierme R. Factor VIII complex and endothelial damage. *Lancet* 1975; **1**: 1430.
- Mordes DB, Lazarchick J, Colwell JA, Sens DA. Elevated glucose concentrations increase factor VIII:Ag levels in human umbilical vein endothelial cells. *Diabetes* 1983; **32**: 876–878.
- Greaves M, Pickering C, Knight G, Boulton AJM, Ball J, Ward JD, Preston FE. Changes in the factor VIII complex in diabetic ketoacidosis: evidence of endothelial cell damage? *Diabetologia* 1987; **30**: 160–165.
- Leurs PB, van Oerle R, Hamulyak K, Wolffenbuttel. Tissue factor pathway inhibitor activity in patients with IDDM. *Diabetes* 1995; **44**: 80–84.
- Porta N, Townsend C, Clover GM, Nanson M, Alderson AR, McCraw A, Kohner EM. Evidence for functional endothelial cell damage in early diabetic retinopathy. *Diabetologia* 1981; **20**: 597–601.
- Jensen T. Increased plasma concentration of von Willebrand factor in insulin dependent diabetics with incipient nephropathy. *Br Med J* 1989; **298**: 27–28.
- Jensen T, Bjerre-Knudsen J, Feldt-Rasmussen B, Deckert T. Features of endothelial dysfunction in early diabetic nephropathy. *Lancet* 1989; **1**: 461–463.
- Stehouwer CDA, Stroes ESC, Hackeng WHL, Mulder PGH, den Ottolander GJH. von Willebrand factor and

- development of diabetic nephropathy in IDDM. *Diabetes* 1991; **40**: 971–976.
20. Knobl P, Schernthaner G, Schnack C, Pietschmann P, Griesmacher A, Prager R, Muller M. Thrombogenic factors are related to urinary albumin excretion rate in type 1 (insulin-dependent) and type 2 (non-insulin dependent) diabetic patients. *Diabetologia* 1993; **36**: 1045–1050.
21. Zimmermann J, Schramm L, Waner C, Mulzer E, Henrich HA, Langer R, Heidreder E. Hemorheology, plasma protein composition and von Willebrand factor in type 1 diabetic nephropathy. *Clin Nephrol* 1996; **46**: 230–236.
22. Myrup B, Mathiesen ER, Ronn B, Deckert T. Endothelial function and serum lipids in the course of developing microalbuminuria in insulin dependent diabetes mellitus. *Diabetes Res* 1994; **26**: 33–39.
23. Yokoyama H, Jensen JS, Myrup B, Mathiesen ER, Ronn B, Deckert T. Raised serum sialic acid concentration precedes onset of microalbuminuria in IDDM. A 10-year follow-up study. *Diabetes Care* 1996; **19**: 435–450.
24. Knobl P, Schernthaner G, Schnack C, Pietschmann P, Griesmacher A, Prager R, Muller M. Thrombogenic factors are related to urinary albumin excretion rate in type 1 (insulin-dependent) and type 2 (non-insulin dependent) diabetic patients. *Diabetologia* 1993; **36**: 1045–1050.
25. Plater ME, Ford I, Dent MT, Preston FE, Ward JD. Elevated von Willebrand factor antigen predicts deterioration in diabetic peripheral nerve function. *Diabetologia* 1996; **39**: 336–343.
26. Stehouwer CDA, Zellenrath P, Polak BCP, Baarsma GS, Nauta JJP, Donker AJM, den Ottolander GJH. Von Willebrand factor and early diabetic retinopathy: no evidence for a relationship in patients with type 1 (insulin dependent) diabetes mellitus and normal urinary albumin excretion. *Diabetologia* 1992; **35**: 555–559.
27. Stehouwer CDA. Is the increase in plasma von Willebrand factor concentrations in diabetes mellitus related to insulin levels? *Thromb Haemostas* 1994; **71**: 398–399.
28. Yaqoob M, Patrick AW, McClelland P, Stevenson A, Mason H, White MC, Bell GM. Relationship between markers of endothelial dysfunction, oxidant injury and tubular damage in patients with insulin dependent diabetes mellitus. *Clin Sci* 1993; **85**: 557–562.
29. Stehouwer CDA, Fischer HRA, van Kuijk AWR, Polak BCP, Donker AJM. Endothelial dysfunction precedes development of microalbuminuria in IDDM. *Diabetes* 1995; **44**: 561–564.
30. Earle KA, Mishra M, Morocutti A, Barnes D, Stephens E, Chambers J, Viberti GC. Microalbuminuria as a marker of silent myocardial ischaemia in IDDM patients. *Diabetologia* 1996; **39**: 854–856.
31. Stehouwer CDA, Donker AJM. Urinary albumin excretion and cardiovascular disease risk in diabetes mellitus: is endothelial dysfunction the missing link? *J Nephrol* 1993; **6**: 72–92.
32. Deckert T, Yokoyama H, Mathiesen E, Ronn B, Jensen T, Feldt-Rasmussen B, Borch-Johnsen K, Jensen JS. Cohort study of predictive value of urinary albumin excretion for atherosclerotic vascular disease in patients with insulin dependent diabetes. *Br Med J* 1996; **312**: 871–874.
33. Jansson JH, Nilsson TK, Johnson O. von Willebrand factor in plasma: a novel risk factor for recurrent myocardial infarction and death. *Br Heart J* 1991; **66**: 351–355.
34. Thompson SG, Kienast J, Pyke SDM, Haverkate F, van de Loo J, et al. Hemostatic factors and the risk of myocardial infarction or sudden death in patients with angina pectoris. *New Engl J Med* 1995; **332**: 635–641.
35. Jones SL, Close CF, Mattock MB, Jarrett RJ, Keen H, Viberti GC. Plasma lipid and coagulation factor concentrations in insulin dependent diabetics with microalbuminuria. *Br Med J* 1989; **298**: 487–490.
36. Boffa MC. Considering cellular thrombomodulin distribution and its modulating factors can facilitate the use of plasma thrombomodulin as a reliable endothelial marker. *Haemostasis* 1996; **26** (suppl 4): 233–243.
37. Galajda P, Martinka E, Mokan M, Kubisz P. Endothelial cell markers in diabetes mellitus. *Thromb Res* 1997; **85**: 63–65.
38. Iwashima Y, Sato T, Watanabe K, Ooshima E, Hiraishi S, Ishii H, Kazama M, Makino I. Elevation of plasma thrombomodulin level in diabetic patients with early diabetic nephropathy. *Diabetes* 1990; **39**: 983–988.
39. Gruden G, Pagano G, Romagnoli R, Frezet D, Olivetti C, Cavallo-Perin P. Thrombomodulin levels in insulin dependent diabetic patients with microalbuminuria. *Diabetic Med* 1995; **12**: 258–260.
40. Gabat S, Keller C, Kempe HP, Amiral J, Ziegler R, Ritz E, et al. Plasma thrombomodulin: a marker for microvascular complications in diabetes mellitus. *Vasa* 1996; **25**: 233–241.
41. Leurs PB, van Oerle R, Hamulyak K, Wolffenbuttel BHR. Tissue factor pathway inhibitor activity in patients with IDDM. *Diabetes* 1995; **44**: 80–84.
42. Seigneur M, Dufourcq P, Gin H, Delafaye C, Amiral J, Pruvost A, Boisseau MR. Plasma thrombomodulin levels increase with the severity of the diabetic retinopathy. *Blood Coag Fibrin* 1994; **5**: 845–846.
43. McLaren M, Elhadd TA, Bancroft A, Connacher M, Belch JFF. Markers of endothelial cell damage and oxidative stress in juvenile diabetics (Abstract). *Blood Coag Fibrin* 1997; **8**: 480.
44. Newman W, Beall LD, Carson CW, Hunder GG, Graben N, Randhawa ZI, et al. Soluble E-selectin is found in supernatants of activated endothelial cells and is elevated in the serum of patients with septic shock. *J Immunol* 1993; **150**: 644–654.
45. Gearing AJH, Hemingway I, Pigott R, Hughes J, Rees AJ, Cashman SJ. Soluble forms of vascular adhesion molecules, E-selectin, ICAM-1, and VCAM-1; pathological significance. *Annals NY Acad Sci* 1992; **667**: 324–331.
46. Steiner M, Reinhardt KM, Krammer B, Ernst B, Blann AD. Increased levels of soluble adhesion molecules in type 2 (non-insulin dependent) diabetes mellitus are independent of glycaemic control. *Thromb Haemostas* 1994; **72**: 979–984.
47. Cominiacini L, Pasini AF, Garbin U, Davoli A, De Santis A, Campagnola M et al. Elevated levels of soluble E-selectin in patients with IDDM and NIDDM: relation to metabolic control. *Diabetologia* 1995; **38**: 1122–1124.
48. Fasching P, Veitl M, Rohac M, Strelci C, Schneider B, Waldhausl W, Wagner OF. Elevated concentrations of circulating adhesion molecules and their association with microvascular complications in insulin dependent diabetes mellitus. *J Clin Endocrin Metabol* 1996; **81**: 4313–4317.
49. Blann AD, McCollum CN. Circulating endothelial cell/leukocyte adhesion molecules in atherosclerosis. *Thromb Haemostas* 1994; **72**: 151–154.
50. Kirk G, Elhadd TA, Robb R, McLaren M, Stonebridge P, Belch JFF. Soluble E-selectin levels in diabetic patients with severe peripheral arterial occlusive disease (Abstract). *Blood Coag Fibrin*. 1996; **7**: 2.
51. Kirk G, Elhadd T, McLaren M, Greene S, Belch JFF.



- Dietary essential fatty acid supplementation in young patients with insulin dependent diabetes mellitus attenuates endothelial activation and/or damage (Abstract). *Blood Coag Fibrin* 1997; **8**: 467.
52. Lerman A, Edwards BS, Hallett JK, Heublein DM, Sandberg SM, Burnett JC. Circulating and tissue endothelin immunoreactivity in advanced atherosclerosis. *New Engl J Med* 1991; **325**: 997–1001.
  53. Takahashi K, Gbatei MA, Lam HC, O'Halloran DJ, Bloom SR. Elevated plasma endothelin in patients with diabetes mellitus. *Diabetologia* 1990; **33**: 306–310.
  54. Collier A, Leach JP, McLennan A, Jardine A, Morton JJ, Small M. Plasma endothelin-like immunoreactive levels in IDDM patients with microalbuminuria. *Diabetes Care* 1992; **15**: 1038–1040.
  55. Kawamura M, Ohgawara H, Naruse M, Suzuki N, Iwasaki N, Naruse K, *et al.* Increased plasma endothelin in NIDDM patients with retinopathy. *Diabetes Care* 1992; **15**: 1396–1397.
  56. Malamitisi-Puchner A, Economou E, Katsouyanni K, Krachaliou F, Delis D, Bartsocas CS. Endothelin 1–21 plasma concentrations in children and adolescents with IDDM. *J Ped Endocrin Metabol* 1996; **9**: 463–468.
  57. Smulders RA, Stehouwer CD, Olthof CG, van Kamp GJ, Teerlink T, de Vries PM, Donker AJ. Plasma endothelin levels and vascular effects of intravenous L-arginine infusion in subjects with uncomplicated insulin dependent diabetes mellitus. *Clin Science* 1994; **87**: 37–43.
  58. Properzi G, Terenghi G, Gu XH, Poccia G, Pasqua R, Francavilla S, Polak JM. Early increase precedes a depletion of endothelin-1 but not of von Willebrand factor in cutaneous microvessels of diabetic patients. A quantitative immunohistochemistry study. *J Pathol* 1995; **175**: 243–252.
  59. Kirilov G, Dakovska L, Borisova AM, Krivoshiev S, Nentchev N. Increased plasma endothelin levels in patients with IDDM and end stage vascular complications. *Hormone & Metabol Res* 1994; **26**: 119–120.
  60. Patino R, Ibarra J, Molino A, Fernandez-Durango R, Moya J, Fernandez-Cruz A. Increased plasma endothelin in diabetes: an atherosclerosis marker? *Diabetologia* 1994; **37**: 333–334.
  61. Gruden G, Cavallo-Perin P, Bazzan M, Stella S, Vuolo A, Pagano G. PAI-1 and factor VII activity are higher in IDDM patients with microalbuminuria. *Diabetes* 1994; **43**: 426–429.
  62. Lindahl AK, Sandset PM, Abildgaard U. The present status of tissue factor pathway inhibitor. *Blood Coag Fibrinolys* 1992; **3**: 439–449.
  63. Sandset PM, Bendz B. Tissue factor pathway inhibitor: Clinical deficiency states. *Thromb Haemost* 1997; **78**: 467–470.
  64. Bajaj MS, Bajaj SP. Tissue factor pathway inhibitor: Potential therapeutic applications. *Thromb Haemost* 1997; **78**: 471–477.
  65. Yokoyama H, Myrup B, Rossing P, Ostergaard PB. Increased tissue factor pathway inhibitor activity in IDDM patients with nephropathy. *Diabetes Care* 1996; **19**: 441–445.
  66. Skrha J, Hodinar A, Kvasnicka J, Stibor V, Sperl M, Stolba P, Hilgertova J. Early changes of serum N-acetyl beta glucosaminidase, tissue plasminogen activator and erythrocyte superoxide dismutase in relation to retinopathy in type 1 diabetes mellitus. *Clin Chem Acta* 1994; **229**: 5–14.
  67. de Bono D. Significance of raised plasma concentrations of tissue-type plasminogen activator and plasminogen activator inhibitor in patients at risk from ischaemic heart disease. *Br Heart J* 1994; **71**: 504–507.
  68. Hekman CM, Loskutoff DJ. Fibrinolytic pathways and the endothelium. *Seminars Thromb Haemost* 1987; **13**: 514–527.
  69. Erikson LA, Ginsberg MN, Loskutoff DJ. Detection and partial characterisation of an inhibitor of plasminogen activator in human platelets. *J Clin Invest* 1984; **74**: 1465–1472.
  70. Wang DL, Pan YT, Wang JJ, Cheng CH, Liu CY. Demonstration of a functionally active tPA-like plasminogen activator in human platelets. *Thromb Haemostas* 1994; **71**: 493–498.
  71. Allen RA, Kluft C, Brommer EJP. The acute effect of smoking on fibrinolysis: increase in the activity level of circulating extrinsic (tissue-type) plasminogen activator. *Eur J Clin Invest* 1984; **14**: 354–361.
  72. Lampeter ER, Hishimoto TK, Rothlein R, Mainolfi EA, Bertrams J, Kolb H, Martin S. Elevated levels of circulating adhesion molecules in IDDM patients and in subjects at risk for IDDM. *Diabetes* 1992; **41**: 1668–1671.
  73. DeLisser HM, Newman PJ, Albelda SM. Molecular and functional aspects of PECAM-1/CD31. *Immunol Today* 1994; **15**: 490–495.
  74. Blann AD, Wadley MS, Dobrotova M, Sanders P, Jayson MIV, McCollum CN. Soluble PECAM in inflammatory vascular disease, atherosclerotic vascular disease, and in cancer. *Blood Coag Fibrin* 1998; in press.
  75. Schleiffenbaum B, Spertini O, Tedder TF. Soluble L-selectin is present in human plasma at high levels and retains functional activity. *J Cell Biol* 1992; **119**: 229–238.
  76. Saika T, Aoki N. Studies of plasma ECMR-III (CD44) and PECAM-1 (CD31) levels in diabetic patients (Abstract). *Diabetologia* 1997; **40** (suppl 1): A401.
  77. Jang Y, Lincoff AM, Plow EF, Topol EJ. Cell adhesion molecules in coronary artery disease. *J Am Coll Cardiol* 1994; **24**: 1591–1601.
  78. Kansas GS. Selectins and their ligands: current components and controversies. *Blood* 1996; **88**: 3259–3287.
  79. Frenette PS, Wagner DD. Adhesion molecules. *New Engl J Med* 1996; **334**: 1526–1529 and **335**: 43–45.
  80. van de Stolpe A, van der Saag PT. Intercellular adhesion molecule-1. *J Mol Med* 1996; **74**: 13–33.
  81. Roep BO, Heidenthal E, de Vries RRP, Kolb H, Martin S. Soluble forms of intercellular adhesion molecule-1 in IDDM. *Lancet* 1994; **343**: 1590–1593.
  82. Uchikoshi F, Ito T, Kamiike W, Moriguchi A, Nozaki S, Ito A, *et al.* Anti-ICAM/LFA-1 monoclonal antibody therapy prevents graft rejection and IDDM recurrence in BB rat pancreas transplantation. *Transpl Proc* 1995; **27**: 1527–1528.
  83. Yagi N, Yokono K, Amano K, Nagata M, Tsukamoto K, Hasegawa Y, *et al.* Expression of intercellular adhesion molecule 1 on pancreatic beta cells accelerates beta cell destruction by cytotoxic t cells in murine autoimmune disease. *Diabetes* 1995; **44**: 744–752.
  84. Accardo-Palumbo A, Triolo G, Giardina E, Carbone MC, Ferrante A, Triolo G. Detection of anti-myeloperoxidase antibodies in the serum of patients with type 1 diabetes mellitus. *Acta Diabetol* 1996; **33**: 103–107.
  85. Esser P, Bresgen M, Fischbach R, Heimann K, Wiedemann P. Intercellular adhesion molecule-1 in plasma and vitreous from patients with vitreoretinal disorders. *German J Ophthalmol* 1995; **4**: 269–274.
  86. Tsukamoto K, Yokono K, Amano K, Nagata M, Yagi N, Tominaga Y, *et al.* *Cell Immunol* 1995; **165**: 193–201.
  87. Abeygunaratne R, Jude E, Young MJ, Bennett J, Abott C, Boulton AJM, Douglas JT. Cellular adhesion molecules in diabetes mellitus: ICAM-1 is a predictor of vascular complications (Abstract). *Thromb Haemostas* 1997; **143** (suppl).



88. Shaw S, Pegrum GD, Wolff S, Ashton WL. Platelet adhesiveness in diabetes mellitus. *J Clin Path* 1967; **20**: 845–847.
89. Colwell JA, Lopes-Virella MF. A review of the development of large vessel disease in diabetes mellitus. *Am J Med* 1988; **85** (suppl 5A): 113–118.
90. Patrick AW, Oliver MD, Howie AF, Dawes J, Macintyre CC, Frier BM. Urinary excretion of beta thromboglobulin and N-acetyl beta-d-glucosaminidase in type 1 diabetes: potential indicators or early nephropathy? *Diabetes Metabol* 1990; **16**: 441–447.
91. Vicari AM, Macagni A. Primary platelet activation in recent onset type 1 diabetes mellitus. *Scand J Clin Lab Invest* 1990; **50**: 429–432.
92. Tschoepe D, Ostermann H, Huebinger A, Ziegler D, Wiefels K, Gries FA. Elevated platelet activation in type 1 diabetes with chronic complications under long-term near-normoglycaemic control. *Haemostasis* 1990; **20**: 93–98.
93. Rak K, Beck P, Udvardy M, Pfliegler G, Misz M, Boda Z. Plasma levels of beta thromboglobulin and factor VIII related antigen in diabetic children and adults. *Thromb Res* 1983; **29**: 155–162.
94. Bayraktar M, Dundar S, Kirazli S, Teletar F. Platelet factor 4, beta thromboglobulin and thrombospondin levels in type I diabetes mellitus patients. *J Int Med Res* 1994; **22**: 90–94.
95. Huszka M, Kaplar M, Rejto L, Tornai I, Palatka K, Laszlo P, Udvardy M. The association of reducing endothelium derived relaxing factor-NO production with endothelial damage and increased in vivo platelet activation in patients with diabetes mellitus. *Thromb Res* 1997; **86**: 173–180.
96. Palumbo PJ, O'Fallon WM, Osmundson PJ, Zimmerman BR, Langworthy AL, Kazmier FJ. Progression of peripheral occlusive arterial disease in diabetes mellitus. What factors are predictive? *Arch Intern Med* 1991; **151**: 717–721.
97. Fijnheer R, Frijns CJM, Korteweg J, Rommes H, Peters JH, Sixma JJ, Nieuwenhuis HK. The origin of P-selectin as a circulating plasma protein. *Thromb Haemostas* 1997; **77**: 1081–1085.
98. Blann AD, Lip GYH. Hypothesis: is soluble P-selectin a marker of platelet activation? *Atherosclerosis* 1997; **128**: 135–138.
99. Blann AD, Lip GYH, Beevers DG, McCollum CN. Soluble P-selectin: a comparison with platelet and endothelial cell markers. *Thromb Haemostas* 1997; **77**: 1077–1080.
100. Jilma B, Fasching P, Ruthner C, Rumplmayr A, Ruzicka S, Kapiotis S, et al. Elevated circulating P-selectin in insulin dependent diabetes mellitus. *Thromb Haemost* 1996; **76**: 328–332.
101. Rauch U, Schwippert B, Schultheiss HP, Tschoepe D. Platelet activation is dependent on microangiopathic complications in insulin dependent diabetes mellitus (Abstract). *Thromb Haemostas* 1997; (suppl): 323.
102. Tschoepe D, Driesch E, Schwippert B, Lampeter EF. Activated platelets in subjects at increased risk of IDDM. DENIS Study Group. *Diabetologia* 1997; **40**: 573–577.
103. Schror K. Blood vessel wall interaction in diabetes. *Diabetes* 1997; **46** (suppl 2): S115–S118.
104. Johnstone MT, Creager SJ, Scales KM, Cusco JA, Lee BK, Creager MA. Impaired endothelium dependent vasodilation in patients with insulin dependent diabetes mellitus. *Circulation* 1993; **88**: 2510–2516.
105. Barnett AH. Pathogenesis of diabetic microangiopathy: an overview. *Am J Med* 1991; **90** (suppl 6A): 67s–73s.
106. Stehouwer CDA, Lambert J, Donker AJM, van Hinsbergh VMW. Endothelial dysfunction and pathogenesis of diabetic angiopathy. *Cardiovasc Res* 1997; **34**: 55–68.
107. Gerrity RG, Antonov AS. The pathogenesis of atherosclerosis. *Diabetologia* 1997; **40**: S108–S110.